


Cell Membrane Physiology

reading: p. 126-129, 134-139

Human Cell- Principal Organelles




Review: Table 3.1, pp. 50-52

Plasma Membrane

- Separates intracellular & extra-cellular spaces
- Regulates entry & exit of substances
- Referred to as the Fluid Mosaic Model

Cell Membrane – Fluid-Mosaic Model



Membrane Function: Transport

- Selective permeability
 - 1) *Passive transport* (no energy)
 - simple diffusion, via gradients
 - facilitated diffusion, via carriers
 - 2) *Active transport* (energy)
 - 3) *Bulk transport*


Passive Transport: Simple Diffusion

Need:

- Concentration gradient
- Permeability to the diffusing substance



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Solute Diffusion



Gradients

Types: concentration, electrochemical, pressure, thermal



Rate of diffusion depends upon:

1) Magnitude of concentration gradient

- Driving force of diffusion

2) Permeability of the membrane

- Neuronal cell membrane 20x more permeable to K^+ than Na^+

3) Temperature


- Higher temperature, faster diffusion rate

4) Surface area of the membrane

- Microvilli increase surface area

Passive Transport:

“Carrier-mediated”



Active Transport

- 1) movement of molecules and ions against their concentration gradients**
- 2) uses specific protein carriers**
- 3) requires ATP**
- 4) 2 types: primary, secondary**


Primary Active Transport

1) Molecule or ion binds to carrier site

2) Binding stimulates breakdown (hydrolysis) of ATP


3) Conformational change moves molecule to other side of membrane.

Thus, direct use of ATP



Primary Active Transport: The Na⁺ - K⁺ Pump

- ◆ An *exchange* pump
- ◆ 3 Na⁺ out for every 2 K⁺ in
- ◆ Energy is used to move ions against their gradients




Secondary Active Transport

- ◆ Coupled transport
- ◆ Energy for uphill movement of glucose is obtained from transport of Na^+ down its [gradient]

Thus, indirect use of ATP
(to maintain Na^+ gradient)

- ◆ Cotransport
- ◆ Countertransport

Cotransport of Na^+ and Glucose into a Cell




Bulk Transport

- Many large molecules moved at the same time

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Endocytosis




Invagination

Formation of pouch

Formation of vesicle

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Exocytosis



Joining of vesicle with plasma membrane

Secretion of cellular product







Figure 3.4, p. 54